

# REHABILITATION GUIDELINES ULNAR/RADIAL COLLATERAL LIGAMENT REPAIR

### Immediate Post Op

DOS:

-Pt. released from surgery in thumb spica cast or splint, with surgical pins in place.

## 2 WEEKS Post-op

Begin Date:

- cast or rigid splint applied

# 6 WEEKS Post-op

Begin Date:

- cast & pins removed
- referred to PT/OT for gamekeeper's thumb splint
- -AROM to thumb, hand, wrist; if necessary, forearm, elbow, shoulder
- PROM to wrist
- after 24 hrs, pt. may begin scar remodel techniques

#### 8 WEEKS Post-op

Begin Date:

- Putty exercises; instruct pt. on proper technique to avoid stress to UCL or RCL (see exercise sheet, next page)
- PRE for wrist, hand, and UE

#### 10 WEEKS Post-op

Begin Date:

- Patient released from splint (unless activities place high stress on thumb joint—gymnastics, carpentry, skiing; these patients should continue splint use)
- Return to normal activities if WNL for grip, pinch, and OK from surgeon

#### 16 to 18 WEEKS Post-op

Begin Date:

- repair to ligament is now mature
- Activities that stress the UCL now ok without splint, per surgeon's release—may require figure 8 taping, etc.

#### **EXERCISES FOR UCL LIGAMENT REPAIR**

ıy.
1

#### THUMB EXERCISES

Position: Face table. Both palms are flat on the table

- 1. Slide the thumb in toward the index finger; then slide away from index finger
- 2. Raise thumb up toward ceiling; hold seconds.

Position: Face table. Palm facing toward ceiling.

- 1. Bring thumb in toward index finger; then, raise thumb towards ceiling (make an "L" with your hand and thumb).
- 2. Blocking: stabilize each portion of the thumb while bending/straightening each joint individually. (MPJ, IPJ).
- 3. Opposition: reach for the little finger with the thumb tip.

#### WRIST EXERCISES

Position: forearm resting on table, wrist and hand hanging off table edge.

- 1. Palm down: make a fist. Reach toward the floor, then straighten fingers and reach to ceiling.
- 2. Palm up: Reach for floor, then reach for the ceiling.
- 3. Thumb up: reach for floor, then reach for ceiling.

Position: Sit or stand, away from table

- 1. Put palms together in "prayer" position; lift elbows for wrist stretch
- 2. Put backs of hands together, fingers relaxed toward floor: bring elbows down for wrist stretch.

Position: Sit or stand with elbows touching ribs, close to your side.

1. Move palm up toward ceiling, then turn palm down toward floor.

#### ADDITIONAL EXERCISES FOR STIFF ARM:

Bend and straighten elbow.

Thumb up, raise arm up straight up.

Scar massage: Massage with lotion on or around scar

Desensitization: Begin with light pressure and soft fabric. Progress to moderate pressure and various textures, rubbing along scar or sensitive area.